



Plymouth County Outreach Project

PAMELA KELLEY

KELLEY RESEARCH ASSOCIATES


STONEHILL COLLEGE

SEAN VARANO

KELLEY RESEARCH ASSOCIATES

ROGER WILLIAMS UNIVERSITY

AUGUST 4, 2020




County-wide Preliminary Data: 1/1/20 – 7/31/20

Type of Incident: 1/1/20 – 7/31/20



Kelley Research Associates




There were a total of **838 incidents (OD and At-Risk)** recorded in Plymouth County between 1/1/20 and 7/31/20: **81 (10%) were fatal; 651 (78%) were non-fatal;** and 106 (12%) were categorized as “at-risk” individuals.

Fatal Incident Monthly Trend



There was on average 11.6 fatal ODs per month, or 81 total, in the County between 1/1/20 and 7/31/20. This is a 5% decrease over the 2019 monthly average of 12.2 or 85 total, for the same time frame.


Non-Fatal Incident Monthly Trend



Kelley Research Associates


There was on average **93 non-fatal ODs per month, or 651 total**, in the County between 1/1/20 and 7/31/20. This is a **<1% decrease** over the 2019 monthly average of 93.4, or 654 total, for the same time frame. Note, there was a 23% increase in non-fatal overdoses between June 2020 and July 2020.

At-Risk Monthly Trend



There was on average **15.1 At-Risk entries per month, or 106 total**, in the County between 1/1/20 and 7/31/20. This is an **3% increase** over the 2019 monthly average of 14.7, or 103 total, for the same time frame.

Incident vs. Resident Town: 1/1/20 – 7/31/20




Kelley Research Associates

Analyzing just the 732 overdose incidents, **508 (69%)** of the OD incidents between 1/1/20 and 7/31/20 occurred in same town that the person lived in. However, in **224 (31%)** of the incidents the person was from a different town than where the incident occurred.

Naloxone “Saves”: 1/1/20 – 7/31/20

Twenty percent (143) of these incidents involved Naloxone being administered by a 3rd party family/friend prior to first responders arriving.




■ Save
■ Fatal

Kelley Research Associates

Again, analyzing just the 732 overdose incidents, Naloxone was administered in **504 (69%)** of the reported OD incidents. **Ninety-four percent (473)** of these events were non-fatal, while **6% (31)** of these overdoses resulted in a fatality.

Follow-Up Visit Contacts 1/1/20 – 7/31/20

- **514** attempted Follow-up Contacts
- **53% (274)** were successful in having contact with either the individual who overdosed, a family/friend, or both.




Fifty-four percent (147) of the successful follow-up visits (274) resulted in contact with the individual that had overdosed, **35% (97)** resulted in contact with Family/Friends, and **11% (30)** resulted in contact with both the individual and family/friends.

Successful SUD Contact Outcomes

1/1/20 – 7/31/20

Kelley Research Associates




Of the 177 successful contacts with the person the overdosed (SUD Only and Both): **67% (119)** resulted in the individual accepting treatment from the outreach team; **14% (25)** declined the treatment options; **14% (24)** of individuals were already seeking treatment; and **5% (9)** of the follow-ups did not have a response to these measures recorded in the system.

Tier 2 Follow-Up Analysis

Tier 2: Follow-Up Visit Contacts 1/1/20 – 7/31/20


- **322** attempted outreach visits (in-person, phone, electronic, other)
- **63% (202)** were successful in having contact with either the individual who overdosed, a family/friend, or both.



Seventy-three percent (148) of the successful follow-up visits (202) resulted in contact with the Tier 2 client, **17% (35)** resulted in contact with Family/Friends, and **10% (19)** resulted in contact with both the Tier 2 client and family/friends.

Tier 2: Successful SUD Contact Outcomes

1/1/20 – 7/31/20



Of the 167 successful contacts with the Tier 2 client (SUD Only and Both): **30% (50)** resulted in the individual accepting treatment from the outreach team; **3% (5)** declined the treatment options; **59% (99)** of individuals was already seeking treatment; and **8% (13)** of the follow-ups did not have a response to these measures recorded in the system.